



Alloantigenic recognition properties of CD8⁺ regulatory T cells

Elodie Picarda, Ignacio Anegon, Carole Guillonnet

► To cite this version:

Elodie Picarda, Ignacio Anegon, Carole Guillonnet. Alloantigenic recognition properties of CD8⁺ regulatory T cells. 6th european workshop on immune-mediated inflammatory diseases, Nice, France. pp.P29. inserm-00643968

HAL Id: inserm-00643968

<https://www.hal.inserm.fr/inserm-00643968>

Submitted on 23 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Alloantigenic recognition properties of CD8⁺ regulatory T cells

Elodie Picarda*, Ignacio Anegón, Carole Guillonnet

From 6th European Workshop on Immune-Mediated Inflammatory Diseases
Nice, France. 23-25 November 2011

Background

We recently reported that in a rat major histocompatibility complex (MHC) mismatched heart allograft model, treatment with CD40Ig, a chimeric molecule that blocks CD40L, leads to indefinite allograft survival mediated by CD8⁺CD45RC^{low} Tregs [1]. Although essential, the exact role of TCR/MHC/peptide interaction in Treg activity is still unknown. We therefore characterize the allogeneic peptide(s) recognized and the TCR usage of the CD8⁺CD45RC^{low} Tregs.

Material and methods

Allogeneic peptide(s) were derived from polymorphic regions of donor MHC molecules [2,3]. Sixty-two overlapping peptides of sixteen amino acids (aa) were tested in a coculture of Tregs with syngeneic pDCs (ratio 4:1). Moreover, the repertoire of the TCR of the CD8⁺ Tregs was studied by flow cytometry analysis and sequencing the CDR3 region.

Results

After six days of culture, two peptides in particular led to the activation of Tregs, as shown by the upregulation of the CD25 molecule (from 25.89% to 29.27% of CD25 expression). These activator peptides were characterized by prominent amino acids (aa) at rather central position, which could result in a large TCR repertoire diversity of the specific Tregs. We showed previously that CD8⁺CD45RC^{low} Tregs expressed a specific altered V 11 repertoire, with the same CDR3 length in all animals (9 aa). This upregulation was confirmed at the protein level, since 19.9±3.7% of Tregs from a CD40Ig-treated animal expressed the V 11 chain compared to 6.1±2.3% in naïve ones. Sequencing of 160 clones of V 11 TCRs from six long-surviving animals suggested a preferential use of a

aa long CDR3 and a particular J region (J 1.6). Interestingly, conserved sequences were frequently found but no common clonotype was shared between animals, suggesting the private nature of the repertoire.

Conclusion

This study demonstrated that CD8⁺CD45RC^{low} Tregs recognize two potential allogeneic epitopes leading to a private TCR repertoire.

Published: 23 November 2011

References

1. Guillonnet C, Hill M, Hubert FX, Chiffolleau E, Herve C, Li XL, Heslan M, Usal C, Tesson L, Menoret S, et al: CD40Ig treatment results in allograft acceptance mediated by CD8CD45RC T cells, IFN-gamma, and indoleamine 2,3-dioxygenase. *J Clin Invest* 2007, **117**:1096-1106.
2. van Denderen B, Peche H, Gagne K, Usal C, Cuturi MC, Souillou JP: Identification of immunodominant donor MHC peptides following rejection and donor strain transfusion-induced tolerance of heart allografts in adult rats. *Eur J Immunol* 2001, **31**:1333-1339.
3. Ballet C, Renaudin K, Degauque N, Mai HL, Boeffard F, Lair D, Berthelot L, Feng C, Smit H, Usal C, et al: Indirect CD4+ TH1 response, antidonor antibodies and diffuse C4d graft deposits in long-term recipients conditioned by donor antigens priming. *Am J Transplant* 2009, **9**:697-708.

doi:10.1186/1479-5876-9-S2-P29

Cite this article as: Picarda et al: Alloantigenic recognition properties of CD8⁺ regulatory T cells. *Journal of Translational Medicine* 2011 **9**(Suppl 2):P29.